

SAN FRANCISCO DUSKY-FOOTED WOODRAT (*NEOTOMA FUSCIPES ANNECTENS*)

DFG: Species of Special Concern



Description

The San Francisco woodrat, *Neotoma fuscipes annectens*, is one of eleven described subspecies of the dusky-footed woodrat (Hooper 1938) found in forest and shrubland communities throughout much of California, the Willamette Valley of Western Oregon, and northern Baja California (Carraway and Verts 1991). Dusky-footed woodrats are medium-sized rodents ranging from 200-400g. The head and body length range from 7 3/5 to 9 inches (193-229 mm) long with a tail length of 6 4/5 to 8 2/3 inches (173-220 mm). The body coloring is brown/grey with white/grey underside and white/dusky coloring on feet. The woodrats have a hairy brown trail usually with a lighter underside, and large ears (Burt and Gossenheider 1980).

Range/distribution The San Francisco woodrat can be found throughout the San Francisco Bay area in grasslands, scrub and wooded areas (Hall 1981).

Life history and habitat

Woodrats are, for the most part, generalist herbivores. They consume a wide variety of nuts and fruits, fungi, foliage and some forbs (Linsdale and Tevis 1951). Many species are good climbers and rock dwellers, and dusky-footed woodrats are highly arboreal (Kelly 1990). Evergreen or live oaks and other thick-leaved trees and shrubs are important habitat components for *N. fuscipes*. (Kelly 1990, Williams et al. 1992).

Dusky-footed woodrats are well known for their large terrestrial stick houses, some of which can last for twenty or more years (Linsdale and Tevis, 1951). At the Hastings Reserve, Monterey County, dusky-footed woodrat houses range from 60 cm (2 feet) to 150 cm (5 feet) in height, and can be 120 cm (4 feet) to 240 cm (8 feet) in basal diameter. Houses typically are placed on the ground against or straddling a log or exposed roots of a standing tree, and, are often located in dense brush. Nests are also placed in the crotches and cavities of trees and in hollow logs. Sometimes arboreal nests are constructed but this behavior seems to be more common in habitat with evergreen trees such as live oak.

Photo: Photo of a dusky-footed woodrat by CSUS/ESRP

References

- Burt, W.H. and R.P. Gossenheider 1980. Peterson field guide to mammals . Houghton Mifflin Company, 289pp.
- Carraway, L. N., and B. J. Verts. 1991. *Neotoma fuscipes* . Mammalian Species 386:1-10.
- Hall, E.R. 1981. Mammals of north america . John Wiley and Sons, 1181 pp.
- Hooper, E.T. 1938. Geographical variation in wood rats of the species *Neotoma fuscipes* . Univ. Calif. Publ. Zool. , 42, 213-246.
- Kelly, P. A. 1990. Population ecology and social organization of dusky-footed woodrats, *Neotoma fuscipes* . Ph.D. dissertation, Univ. of California, Berkeley, 191 pp.
- Linsdale, J. M., and L. P. Tevis, Jr. 1951. The dusky-footed wood rat . Univ. California Press, Berkeley and Los Angeles, 664 pp.
- National Parks Service and Golden Gate National Parks Association. Fort baker plan. URL: <http://www.nps.gov/goga/ftbaker/eis/Appendices.pdf>
- Williams, D.F. 1986. Mammalian species of special concern in California . California Department of Fish and Game (pp. 45-46).
- Williams, D. F. 1993. Population censuses of riparian brush rabbits and riparian woodrats at Caswell Memorial State Park during January 1993. Final Report to California Dept. Parks and Recreation, Lodi, CA , 15 pp.
- Williams, D.F., J. Verner, H.F. Sakai, and J.R. Waters. 1992. General biology of major prey species of the California spotted owl. USDA Forest Service Gen. Tech. Rep., PSW-GTR-133:207-221.